

VOLTERRA TYPE INTEGRAL OPERATORS AND COMPOSITION OPERATORS ON MODEL SPACES

TESFA MENGESTIE

ABSTRACT. We study some mapping properties of Volterra type integral operators and composition operators on model spaces. We also discuss and give out a couple of interesting open problems in model spaces where any possible solution of the problems can be used to study a number of other operator theoretic related problems in the spaces.

1. INTRODUCTION

We denote by $\mathcal{H}$ the space of holomorphic functions. We define the Volterra type integral operator on $\mathcal{H}$ induced by a homomorphic symbol g by

$$V_g f(z) = \int_0^z f(w)g'(w)dw.$$

Questions about various operator theoretic properties of V_g expressed in terms of function theoretic conditions on g have been a subject of high interest mainly after the works of Pommerenke [22] and later by Aleman, Cima and Siskakis [4, 6, 7]. In [6], it was proved that V_g on the Hardy space $H^p(\mathbb{D})$, $1 \leq p < \infty$, is bounded if and only if g belongs to the space of $BMOA$, while compactness is characterized in terms of the corresponding $VMOA$ spaces.

In this paper we plan to study the bounded, compact and Hilbert–Schmidt properties of the Volterra type integral operators and composition operators acting between the model spaces K_I^2 and the Hardy space $H^2(\mathbb{C}^+)$ of the upper-half plane. We recall that the model spaces K_I^2 , where I is an inner function, are the subspace of $H^2(\mathbb{C}^+)$ defined by $K_I^2 = H^2(\mathbb{C}^+) \ominus IH^2(\mathbb{C}^+)$. The space K_I^2 is a reproducing kernel Hilbert space with kernel function

$$K_w(z) = \frac{i}{2\pi} \cdot \frac{1 - \overline{I(w)}I(z)}{z - \overline{w}}, \quad k_w(z) = K_w(z) / \|K_w\|_2.$$

Each inner function I has either one or infinitely many components. We call it a one-component if there exists a δ in $(0, 1)$ for which the level set $\{z \in \mathbb{C}^+ : |I(z)| < \delta\}$ is connected. For model spaces generated by such class of inner functions, we prove the following.

Theorem 1.1. *Let I be a one-component inner function on $\mathbb{C}^+$ and g be a holomorphic function on $\mathbb{C}^+$. Then $V_g : K_I^2 \rightarrow H^2(\mathbb{C}^+)$ is*

2010 *Mathematics Subject Classification.* Primary 47B32, 30H20; Secondary 46E22, 46E20, 47B33

Key words and phrases. Volterra type integral operator, bounded, compact, composition operator, model spaces.

(i) *bounded if and only if*

$$\sup_{w \in \mathbb{C}^+} \int_{\mathbb{C}^+} |k_w(z)|^2 |g'(z)|^2 \Im z dA(z) < \infty; \quad (1.1)$$

(ii) *compact if and only if*

$$\lim_{|w| \rightarrow \infty} \int_{\mathbb{C}^+} |k_w(z)|^2 |g'(z)|^2 \Im z dA(z) = 0; \quad (1.2)$$

(iii) *in the Hilbert–Schmidt class $\mathcal{S}_2(K_I^2, H^2(\mathbb{C}^+))$ if and only if*

$$\int_{\mathbb{C}^+} |g'(z)|^2 (1 - |I(z)|^2) dA(z) < \infty, \quad (1.3)$$

where dA denotes the usual Lebesgue area measure.

The criteria found above classify the bounded and compact V_g on Model spaces in terms of their action on the reproducing kernels. As will be seen in Section 4, by converting the boundary integral in Hardy spaces to an area integral in terms of derivatives, we could reduce the boundedness and compactness problems of V_g into questions of bounded and compact Carleson embedding maps (measures) for K_I^2 . Such measures have been identified when I belongs to the class of one-component inner functions: for example see [1, 2, 11, 12, 13, 26]. Despite several attempts made by many experts in the subject, a complete identification of the measures remains open when I belongs to the class of infinitely many component inner functions. For special class of model spaces which admit reproducing kernel Riesz bases corresponding to sparse sequence of points, which includes the de Branges spaces, the author together with K. Seip and Y. Belov [8] have described the measures in terms of a condition analogous to the A_2 weight condition. In what follows we will use those results in [8] to study the bounded and compact properties of V_g . We note in passing that those spaces constitute model spaces generated by the class of infinitely many component inner functions.

1.1. The Volterra type integral operators on $\mathcal{H}(\Gamma, v)$. Let $\Gamma = (\gamma_n)$ be a sequence of distinct complex numbers and $v = (v_n)$ be a weight sequence that satisfies the admissibility condition

$$\sum_{n=1}^{\infty} \frac{v_n}{1 + |\gamma_n|^2} < \infty. \quad (1.4)$$

Any such pair (Γ, v) parameterizes the space $\mathcal{H}(\Gamma, v)$ which consists of all functions

$$f(z) = \sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n} \quad (1.5)$$

for which

$$\|f\|_{\mathcal{H}(\Gamma, v)} = \|(a_n)\|_{\ell_v^2} < \infty, \quad \ell_v^2 = \{(a_n) : \|a\|_v^2 = \sum_{n=1}^{\infty} |a_n|^2 v_n < \infty\} \quad (1.6)$$

and z belongs to the set

$$(\Gamma, v)^* = \left\{ z \in \mathbb{C} : \sum_{n=1}^{\infty} \frac{v_n}{|z - \gamma_n|^2} < \infty \right\}. \quad (1.7)$$

The equation in (1.5) means that we obtain the value of a function f in $\mathcal{H}(\Gamma, v)$ at a point z in $(\Gamma, v)^*$ by computing the weighted discrete Hilbert transform:

$$(a_n) \mapsto \sum_{n=1}^{\infty} \frac{a_n v_n}{z - \gamma_n}, \quad (1.8)$$

which is well defined whenever (a_n) belongs to ℓ_v^2 . This follows from an application of Cauchy–Schwarz inequality along with (1.6) and (1.7). The de Branges spaces, model subspaces of the Hardy space H^2 which admit Riesz bases of reproducing kernels and the Fock-type spaces studied in [10] are all examples of spaces of the kind $\mathcal{H}(\Gamma, v)$. We refer to [8, 9, 18] for detailed accounts. The Carleson measures on $\mathcal{H}(\Gamma, v)$ have been described in [8] when Γ grows at least exponentially, i. e., when

$$\inf_{n \geq 1} \frac{|\gamma_{n+1}|}{|\gamma_n|} > 1. \quad (1.9)$$

When we consider such sparse sequence Γ , it is natural to partition $\mathbb{C}^+$ in the following way. We set $\Omega_1 = \{z \in \mathbb{C}^+ : |z| < (|\gamma_1| + |\gamma_2|)/2\}$ and then for $n \geq 2$

$$\Omega_n = \{z \in \mathbb{C}^+ : (|\gamma_{n-1}| + |\gamma_n|)/2 \leq |z| < (|\gamma_n| + |\gamma_{n+1}|)/2\}.$$

We now state our first result on $\mathcal{H}(\Gamma, v)$.

Theorem 1.2. *Suppose that the sequence Γ satisfies the sparseness condition (1.9) and $g : (\Gamma, v)^* \rightarrow \mathbb{C}^+$ be homomorphic. Then $V_g : \mathcal{H}(\Gamma, v) \rightarrow H^2(\mathbb{C}^+)$ is*

(i) *bounded if and only if*

$$\sup_{n \geq 1} \int_{\Omega_n} \frac{v_n |g'(z)|^2 \Im z dA(z)}{|z - \gamma_n|^2} < \infty \quad (1.10)$$

and

$$\sup_{n \geq 1} \left(\sum_{l=1}^n v_l \sum_{m=n+1}^{\infty} \int_{\Omega_m} \frac{|g'(z)|^2 \Im z dA(z)}{|z|^2} + \sum_{l=n+1}^{\infty} \frac{v_l}{|\gamma_l|^2} \sum_{m=1}^n \int_{\Omega_m} |g'(z)|^2 \Im z dA(z) \right) < \infty; \quad (1.11)$$

(ii) *compact if and only if the "little oh" counterparts of (1.10) and (1.11) holds;*

(iii) *in the Hilbert–Schmidt class $\mathcal{S}_2(\mathcal{H}(\Gamma, v), H^2(\mathbb{C}^+))$ if and only if*

$$\sum_{n=1}^{\infty} \int_{\Omega_n} \frac{v_n |g'(z)|^2 \Im z dA(z)}{|z - \gamma_n|^2} < \infty \quad (1.12)$$

and

$$\sum_{k=1}^{\infty} v_k \sum_{n=k+1}^{\infty} \int_{\Omega_n} \frac{|g'(z)|^2 \Im z dA(z)}{|z|^2} + \sum_{k=n+1}^{\infty} \frac{v_n}{|\gamma_n|^2} \sum_{k=1}^n \int_{\Omega_k} |g'(z)|^2 \Im z dA(z) < \infty. \quad (1.13)$$

Condition (1.10) of the theorem is a condition about the local behavior of the symbol g , while condition (1.11) deals with its global behavior. Combining the two conditions, we see that (1.10) may be replaced by the stronger global condition

$$\sup_{n \geq 1} \int_{\mathbb{C}^+} \frac{v_n |g'(z)|^2 \Im z dA(z)}{|z - \gamma_n|^2} < \infty.$$

In some special cases, this condition, or the original one (1.10) automatically gives condition (1.11). For example if the weight sequence v_n grows at least exponentially and the numbers $v_n/|\gamma_n|^2$ decay at least exponentially with n . The same conclusion follows if the weight sequence v_n is also summable.

2. THE COMPOSITION OPERATOR ON $\mathcal{H}(\Gamma, v)$

Let ψ be a homomorphic function in a given domain. We denote by C_ψ the composition operator $f \mapsto f \circ \psi$. As a consequence of the Littlewood's Subordination Theorem [15], it has long been known that all composition operators are bounded on all the Hardy spaces H^p of the unit disc where $0 < p \leq \infty$. Although corresponding Hardy spaces of the disc and half-plane are isomorphic, composition operators act differently in the two domains. Unlike the case of $H^p(\mathbb{D})$, not all composition operators are bounded on $H^p(\mathbb{C}^+)$.

The study of compact composition operators on $H^2(\mathbb{D})$ first appeared in H. Schwartz [23] thesis in the late sixties though a complete function theoretic characterization, in terms of the inducing map's Nevanlinna counting function, was obtained by Shapiro later in [24]. Unlike $H^p(\mathbb{D})$ again, it was proved in [17] that there exists no compact composition operator on the Hardy space $H^p(\mathbb{C}^+)$. The work in [24] was continued by Y. Lyubarskii and E. Malinnikova [16] for $C_\psi : K_I^2(\mathbb{D}) \rightarrow H^2(\mathbb{D})$ and a complete extension was made when the generating inner function I is one-component. A more general trace ideal criteria for C_ψ was obtained latter in [5]. The problem remains open when I has infinitely many components.

In this section, we study the bounded, compact and Hilbert–Schmidt properties of composition operators on model spaces when the spaces admit normalized reproducing kernel Riesz bases associated to sparse sequences of points in $\mathbb{C}^+$. We note that such spaces represent some classes of model spaces generated by infinitely many component inner functions.

Theorem 2.1. *Suppose that the sequence Γ satisfies the sparseness condition (1.9) and ψ be a nonconstant analytic function on $\mathbb{C}^+$, and m refers to the usual Lebesgue measure on the real line. Then $C_\psi : \mathcal{H}(\Gamma, v) \rightarrow H^2(\mathbb{C}^+)$ is*

(i) *bounded if and only if*

$$\sup_{n \geq 1} \sup_{y \geq 0} \int_{\psi(\Omega_n)} \frac{v_n dx}{|\psi(x + iy) - \gamma_n|^2} < \infty \quad (2.1)$$

and

$$\sup_{n \geq 1} \left(\sum_{l=1}^n v_l \sum_{k=n+1}^{\infty} \sup_{y \geq 0} \int_{\psi(\Omega_k)} \frac{dx}{|\psi(x+iy)|^2} + \sum_{l=n+1}^{\infty} \frac{v_l}{|\gamma_l|^2} \sup_{y \geq 0} \sum_{k=1}^n m(\psi(\Omega_k)) \right) < \infty, \quad (2.2)$$

where $\psi(\Omega_k) = \{\psi(x+iy) : x+iy \in \Omega_k\}$;

- (ii) compact if and only if the "little oh" counterparts of conditions (2.1) and (2.2) hold;
- (iii) in the Hilbert-Schmidt class $\mathcal{S}_2(\mathcal{H}(\Gamma, v), H^2(\mathbb{C}^+))$ if and only if

$$\sum_{k=1}^{\infty} \sup_{y > 0} \int_{\psi(\Omega_k)} \frac{v_k dx}{|\psi(x+iy) - \gamma_k|^2} < \infty \quad (2.3)$$

and

$$\sum_{k=1}^{\infty} v_k \sum_{j=k+1}^{\infty} \sup_{y > 0} \int_{\psi(\Omega_j)} \frac{dx}{|\psi(x+iy)|^2} + \sum_{k=1}^{\infty} \frac{v_k}{|\gamma_k|^2} \sup_{y > 0} \sum_{j=1}^{k-1} m(\psi(\Omega_j)) < \infty. \quad (2.4)$$

3. OPEN PROBLEMS

3.1. Open problem 1. As will be seen in the next section, the proof of our boundedness and compactness results for both C_ψ and V_g rely mainly on previously obtained Carleson measure results on $\mathcal{H}(\Gamma, v)$. When the sequence of points $\gamma_n = n$ and the weight sequences $v_n \simeq 1$, the space $\mathcal{H}(\Gamma, v)$ becomes the classical Paley-Wiener space for which its Carleson measures have long been identified. When the sequences γ_n grows at least exponentially, the measures are completely described in terms of A_2 type condition in [8]. The case when γ_n grows between these two extreme cases remains an interesting open problem. A solution of this will settle the long-standing open problem of identifying the Carleson measures for Model spaces, of course modulo to the existence of reproducing kernel Riesz basis in Model spaces. The novelty and the core of the approach in [8] has been in turning a number of problems including the Carleson measure problem into questions about different mapping properties of Hilbert transforms in weighted spaces of functions and sequences. To extend the approach, one may need to look into all possible interplays between the smoothness (regularity) of the weight sequence (v_n) and the growth of the sequence $\Gamma = (\gamma_n)$.

3.2. Open problem 2. Both of our results in parts iii) of Theorem 1.2 and Theorem 2.1 deal with the Hilbert-Schmidt properties of the operators V_g and C_ψ on $\mathcal{H}(\Gamma, v)$. A more natural and interesting question is to ask the Schatten $\mathcal{S}_p$ class membership of these maps from $\mathcal{H}(\Gamma, v)$ into $H^2(\mathbb{C}^+)$ for all p in the range $0 < p < \infty$. Dealing with these problems would have been easier had we been known a complete characterization of the $\mathcal{S}_p$ class membership of the imbedding maps induced by Carleson measures for the spaces, which itself is still another open problem. A good starting point for all of these problems could be to consider first the case when the sequences γ_n grows at least exponentially. A very special case of the $\mathcal{S}_p$ membership problem pertaining to the embedding maps may be seen in [19].

A word on notation: the notation $U(z) \lesssim V(z)$ (or equivalently $V(z) \gtrsim U(z)$) means that there is a constant C such that $U(z) \leq CV(z)$ holds for all z in the set of a question. We write $U(z) \simeq V(z)$ if both $U(z) \lesssim V(z)$ and $V(z) \lesssim U(z)$.

4. PROOF OF THE RESULTS

Proof of Theorem 1.1. The key tool in the proofs of the first two results is the Littlewood–Paley description of Hardy spaces in terms of derivative :

$$\|f\|_2^2 = 2 \int_{\mathbb{C}^+} |f'(z)|^2 \Im z dA(z) \quad (4.1)$$

for each f in $H^2(\mathbb{C}^+)$. The formula helps to convert a boundary integral to an area integral in terms of derivatives. Since $(V_g f)'(z) = f(z)g'(z)$, an application of the above identity ensures that for each f in K_I^2 we have

$$\|V_g f\|_2^2 = 2 \int_{\mathbb{C}^+} |f(z)|^2 |g'(z)|^2 \Im z dA(z). \quad (4.2)$$

It follows from this that the boundedness of the operator $V_g : K_I^2 \rightarrow H^2(\mathbb{C}^+)$ is equivalent to saying that the measure μ_g where

$$d\mu_g(z) = |g'(z)|^2 \Im z dA(z) \quad (4.3)$$

is a Carleson measure for K_I^2 . Assuming that I is a one-component inner function, a result of Cohn [12] ensures that the measure property holds if and only if the normalized reproducing kernels of K_I^2 are uniformly embedded into $L^2(\mathbb{C}^+, d\mu_g)$ from which the boundedness part of the result follows.

To prove (ii), first from a standard normal family argument, we have that V_g is compact if and only if $V_g f_n \rightarrow 0$ in $H^2(\mathbb{C}^+)$ for each sequence of functions f_n in K_I^2 such that the sequence is uniformly bounded in norm and converges uniformly to zero on compact subsets of $\mathbb{C}^+$. It means that the compactness of V_g is equivalent to saying that the embedding map from K_I^2 into $L^2(\mu_g)$ is compact. Since I is one-component, then the desired conclusion follows from a result of Cohn [12] again.

To prove part (iii), recall that the operator V_g belongs to the Hilbert–Schmidt class if and only if for any orthonormal basis (e_n) of K_I^2 , the sequence $(\|V_g e_n\|_2^2)$ is summable. It follows by (4.2) that

$$\begin{aligned} \sum_{n=1}^{\infty} \|V_g e_n\|_2^2 &\simeq \sum_{n=1}^{\infty} \int_{\mathbb{C}^+} |e_n(z)|^2 |g'(z)|^2 \Im z dA(z) \\ &= \int_{\mathbb{C}^+} \sum_{n=1}^{\infty} |e_n(z)|^2 |g'(z)|^2 \Im z dA(z). \end{aligned}$$

On the other hand, because of the reproducing property of the kernel and Parseval's identity, we have

$$K_w(z) = \sum_{n=1}^{\infty} \langle K_w, e_n \rangle e_n(z) = \sum_{n=1}^{\infty} e_n(z) \overline{e_n(w)} \text{ and } \|K_w\|_2^2 = \sum_{n=1}^{\infty} |e_n(w)|^2. \quad (4.4)$$

The desired conclusion easily follows since

$$\sum_{n=1}^{\infty} |e_n(z)|^2 = \|K_z\|_2^2 \simeq \frac{1 - |I(z)|^2}{\Im z}. \quad (4.5)$$

Proof of Theorem 1.2. Applying the Littlewood–Paley identity again, the estimate

$$\|V_g f\|_2^2 = 2 \int_{\mathbb{C}^+} |f(z)|^2 |g'(z)|^2 \Im z dA(z) \lesssim \|f\|_{\mathcal{H}(\Gamma, v)}^2 \quad (4.6)$$

holds if and only if the embedding map from $\mathcal{H}(\Gamma, v)$ into $L^2(\mathbb{C}, \mu_g)$ is bounded where μ_g is the nonnegative measure defined by (4.3). Then, the boundedness part of the result immediately follows from an application of Theorem 1.1 in [8].

By a result of Nordgren's [20], a sequence of functions f_n in $\mathcal{H}(\Gamma, v)$ converges weakly to zero if and only if it converges pointwise to zero and $\sup_n \|f_n\|_{\mathcal{H}(\Gamma, v)} < \infty$. It follows that V_ψ is compact if and only if the measure μ_g is a vanishing Carleson measure for $\mathcal{H}(\Gamma, v)$. Then an application of Theorem 1.2 in [18] finishes the proof of the compactness part.

It remains to prove the statement about Hilbert–Schmidt membership. First, we may observe that the reproducing kernel of $\mathcal{H}(\Gamma, v)$ at a point λ in $(\Gamma, v)^*$ is explicitly given by

$$k_\lambda(z) = \sum_{n=1}^{\infty} \frac{v_n}{(\bar{\lambda} - \bar{\gamma}_n)(z - \gamma_n)};$$

this is a direct consequence of the definition of the space $\mathcal{H}(\Gamma, v)$. Furthermore, the sequence $e_n(z) = \sqrt{v_n}/(z - \gamma_n)$ constitutes an orthonormal basis to $\mathcal{H}(\Gamma, v)$. Then

$$\begin{aligned} \|V_g\|_{\mathcal{S}_2}^2 &= \sum_{n=1}^{\infty} \|V_g e_n\|_2^2 \simeq \sum_{n=1}^{\infty} \int_{\mathbb{C}^+} |e_n(z)|^2 |g'(z)|^2 \Im z dA(z) \\ &= \int_{\mathbb{C}^+} \|K_z\|_{\mathcal{H}(\Gamma, v)}^2 |g'(z)|^2 \Im z dA(z) \\ &= \int_{\mathbb{C}^+} \sum_{n=1}^{\infty} \frac{v_n}{|z - \gamma_n|^2} |g'(z)|^2 \Im z dA(z). \end{aligned} \quad (4.7)$$

Using the growth condition (1.9), we observe that if $z \in \Omega_m$, then

$$\sum_{n=1}^{\infty} \frac{v_n}{|z - \gamma_n|^2} \simeq \sum_{n=1}^{m-1} \frac{v_n}{|z|^2} + \frac{v_m}{|z - \gamma_m|^2} + \sum_{n=m+1}^{\infty} \frac{v_n}{|\gamma_n|^2}. \quad (4.8)$$

Plugging this estimate in (4.7), we find that $\|V_g\|_{\mathcal{S}_2}$ is finite if and only if both (1.12) and (1.13) hold.

Proof of Theorem 2.1. Suppose now that ψ is analytic on $\mathbb{C}^+$. Then for each fixed $y > 0$, we define $\psi_y(x) = \psi(x + iy)$ from $\mathbb{R}$ into $\mathbb{C}^+$. We denote by $m\psi_y^{-1}$ the pullback measure

$$m\psi_y^{-1}(E) = m(\psi_y^{-1}(E))$$

for each Borel subset $E \subset \mathbb{C}^+$ where m refers to the usual Lebesgue measure on the real line.

Applying change of variables, we observe

$$\int_{\mathbb{R}} |C_\psi f(x + iy)|^2 dx = \int_{\mathbb{C}^+} |f(z)|^2 dm\psi_y^{-1}(z); \quad (4.9)$$

which holds because of a classical result in measure theory ([14], p.163). This implies that the estimate

$$\|C_\psi f\|_2^2 = \sup_{y>0} \int_{\mathbb{C}^+} |(f \circ \psi)(x + iy)|^2 dx \lesssim \|f\|_{\mathcal{H}(\Gamma, v)}^2 \quad (4.10)$$

holds if and only if $(m\psi_y^{-1})_{y>0}$ constitute a family of Carleson measure for $\mathcal{H}(\Gamma, v)$ with a uniform bound on y in the sense that

$$\sup_{y>0} \frac{\int_{\mathbb{C}^+} |f(z)|^2 dm\psi_y^{-1}(z)}{\|f\|_{\mathcal{H}(\Gamma, v)}^2} < \infty.$$

Then an application of Theorem 1.1 of [8] gives the first part of the result in the theorem.

The compactness part of our result follows easily by similar arguments used to prove the compactness part in Theorem 1.2 along with Theorem 1.2 of [18]. Thus, we remain to prove part (iii) of the result. Proceeding as in the proof of part (iii) of Theorem 1.2 and applying (4.10) we have

$$\begin{aligned} \|C_\psi\|_{\mathcal{S}_2}^2 &= \sum_{n=1}^{\infty} \|C_\psi e_n\|_2^2 = \sum_{n=1}^{\infty} \sup_{y>0} \int_{\mathbb{C}^+} |(e_n \circ \psi)(x + iy)|^2 dx \\ &= \sum_{n=1}^{\infty} \sup_{y>0} \int_{\mathbb{C}^+} |e_n(z)|^2 dm\psi_y^{-1}(z) = \sum_{n=1}^{\infty} \sup_{y>0} \int_{\mathbb{C}^+} \frac{v_n}{|z - \gamma_n|^2} dm\psi_y^{-1}(z) \\ &= \sum_{n=1}^{\infty} \sup_{y>0} \sum_{j=1}^{\infty} \int_{\Omega_j} \frac{v_n}{|z - \gamma_n|^2} dm\psi_y^{-1}(z) \end{aligned} \quad (4.11)$$

Applying the growth condition in (1.9), we observe that if $z \in \Omega_n$, then

$$\sum_{n=1}^{\infty} v_n \sup_{y>0} \sum_{j=1}^{\infty} \int_{\Omega_j} \frac{dm\psi_y^{-1}(z)}{|z - \gamma_n|^2} \simeq \sum_{n=1}^{\infty} v_n \sup_{y>0} \sum_{j=1}^{n-1} \int_{\Omega_j} \frac{dm\psi_y^{-1}(z)}{|z|^2} \quad (4.12)$$

$$+ \sum_{n=1}^{\infty} \sup_{y>0} \left(\int_{\Omega_n} \frac{v_n dm\psi_y^{-1}(z)}{|z - \gamma_n|^2} + \sum_{j=n+1}^{\infty} v_n \int_{\Omega_j} \frac{dm\psi_y^{-1}(z)}{|\gamma_n|^2} \right). \quad (4.13)$$

The sum on the right hand side of (4.12) can be further rewritten as

$$\sum_{n=1}^{\infty} v_n \sup_{y>0} \sum_{j=1}^{n-1} \int_{\Omega_j} \frac{dm\psi_y^{-1}(z)}{|z|^2} = \sum_{n=1}^{\infty} v_n \sup_{y>0} \sum_{j=1}^{n-1} \int_{\psi(\Omega_j)} \frac{dx}{|\psi(x + iy)|^2}. \quad (4.14)$$

On the other hand, the sum on the right hand side of (4.13) becomes

$$\begin{aligned} \sum_{n=1}^{\infty} v_n \sup_{y>0} \left(\int_{\Omega_n} \frac{dm\psi_y^{-1}(z)}{|z - \gamma_n|^2} + \sum_{j=n+1}^{\infty} \int_{\Omega_j} \frac{v_n}{|\gamma_n|^2} dm\psi_y^{-1}(z) \right) \\ \simeq \sum_{n=1}^{\infty} v_n \sup_{y>0} \int_{\psi(\Omega_n)} \frac{dx}{|\psi(x + iy) - \gamma_n|^2} \\ + \sum_{n=1}^{\infty} \frac{v_n}{|\gamma_n|^2} \sup_{y>0} \sum_{j=n+1}^{\infty} m(\psi(\Omega_j)), \end{aligned} \quad (4.15)$$

where $m(\psi(\Omega_j)) = m(\{\psi(x + iy) : x + iy \in \Omega_j\})$. From the above series of estimates, we see that $\|C_\psi\|_{\mathcal{S}_2}$ is finite if and only if the sums in (4.14) and (4.15) are finite, and from which our assertion follows.

REFERENCES

- [1] A. Aleksandrov, *A simple proof of the Volberg–Treil theorem on the embedding of coinvariant subspaces of the shift operator*, Zap. Nauchn. Sem. S. Peterburg Otdel. Mat. Inst. Steklov., 217 (1994), 26–35.
- [2] A. Aleksandrov, *On embedding theorems for coinvariant subspaces of the shift operator II*, J. Math. Sci., 110 (2002), no. 5, 45–64.
- [3] A. Aleman, *A class of integral operators on spaces of analytic functions*, in topics in complex analysis and operator theory, 3–30, Univ. Málaga, Málaga, 2007.
- [4] A. Aleman and J. Cima, *An integral operator on H_p and Hardy’s inequality*, J. Anal. Math. 85 (2001), 157–76.
- [5] A. Aleman, Yu. Lyubarskii, E. Malinnikova, and K.-M. Perfekt, *Trace ideal criteria for embeddings and composition operators on model spaces*, ArXiv: 1307.2652, 2013.
- [6] A. Aleman and A. Siskakis, *An integral operator on H^p* Complex Variables, 28(1995), 149–158.
- [7] A. Aleman and A. Siskaki, *Integration operators on Bergman spaces*, Indiana University Math J., 46(1997), 337–356.
- [8] Y. Belov, T. Mengestie, and K. Seip, *Discrete Hilbert transforms on sparse sequences*, Proc. London Math. Soc. (3) 103 (2011), no. 1, 73–105.
- [9] Y. Belov, T. Mengestie, and K. Seip, *Unitary discrete Hilbert transforms*, J. Anal. Math., 112 (2010), 383–393.
- [10] A. Borichev and Yu. Lyubarskii, *Riesz bases of reproducing kernels in Fock-type spaces*, J. Inst. Math. Jussieu, 9 (2010), 449–461.
- [11] W. Cohn, *Carleson measures and operators on star-invariant subspaces*, J. Oper. Theory, 15 (1986), 181–202.
- [12] W. Cohn, *Carleson measures for functions orthogonal to invariant subspaces*, Pacific J. Math., 103 (1982), 347–364.
- [13] W. Cohn, *Radial embedding theorems for invariant subspaces*, Complex Variables Theory Appl., 17 (1991), 33–42.
- [14] P. Halmos, *Measure Theory*, Van Nostrand, Princeton, NJ, 1950.
- [15] J. Littlewood, *On inequalities in the theory of functions*, Proc. London Math. Soc., 23(1925) 481–519.
- [16] Y. Lyubarskii and E. Malinnikova, *Composition operator on Model spaces*, Recent trends in Analysis: Proceedings of the conference in honor of Nikolai Nikolski, Bordeaux; Theta Foundation, Bucharest, 149–157, 2013.
- [17] V. Matache, *Composition operators on hardy spaces of a half-plane*, Proceedings of the Amer. Math. Soc., 127(1999), 5, 1483–1491.

- [18] T. Mengestie, *Compact Carleson measures from sparse sequences*, Rocky Mountain J. Math., Vol 44, 1 (2014), 223–234.
- [19] T. Mengestie, *Two weight discrete Hilbert transforms and systems of reproducing kernels*, PhD, thesis, Norwegian University of Science and Technology, 2011.
- [20] E. A. Nordgren, *composition operators on Hilbert spaces, lecture notes in math.* springer–Verlag, Berlin, 693(1978), 37–63.
- [21] F. Nazarov and A. Volberg, *The Bellman function, two weighted Hilbert transform, and embeddings of the model space K_θ* , J. Anal. Math., 87(2002), 385–414.
- [22] C. Pommerenke, *Schlichte Funktionen und analytische Funktionen von beschränkter mittlerer Oszillation*, Commentarii Mathematici Helvetici, 52 (1977), no. 4, 591–602.
- [23] H. Schwartz, *Composition Operators on H^p* , Thesis Univ. of Toledo, 1968.
- [24] J. H. Shapiro, *The essential norm of a composition operator*, Annals of Math., 125 (1987), 375–404.
- [25] J. Shapiro and W. Smith, *Hardy spaces that support no compact composition operators*, J. Func. Analysis., 205 (2003), 62–89.
- [26] S. Treil and A. Volberg, *Embedding theorems for invariant subspaces of the inverse shift operator*, translation in J. Soviet Math., 42 (1988), 1562–1572.

DEPARTMENT OF MATHEMATICAL SCIENCES, STORD/Haugesund University College
(HSH), KLINGENBERGVEGEN 8, N-5414 STORD, NORWAY

E-mail address: Tesfa.Mengestie@hsh.no